

Presentation/Publication Information:

A contributed talk at the 8th Foresight Conference on Molecular Nanotechnology in Bethesda, MD, Nov 2-5, 2000. A copy of the talk is attached.

Acknowledgments:

This work is supported under NASA contract DTTS59-99-D-00437 / A61812D.

Curvature Dependent Reactivity of Fullerenes and Nanotubes


Seongjun Park¹, Kyeongjae Cho¹, Deepak Srivastava²

¹Multiscale Simulation Lab., Stanford University

²Nanotechnology at CSC/NAS (IN), NASA Ames Research Center

Curvature Dependent Reactivity

- Example: Hydrogenation of Fullerenes and Nanotubes



- External Surface : Nucleophile
Internal Surface : Electrophile
- Chemical reactivity is a function of the curvature of fullerene.


Nov 5, 2000

MSL, Stanford & NASA Ames

2/20

Dependence on Pyramidalization Angle ?

- The Chemical Reactivity of Fullerene ← The pyramidalization.
- Pyramidalization → Between sp^2 and sp^3
- Between sp^2 and sp^3
 - ↓ Less Energy needed than sp^2
 - ↓ sp^3
- Pyramidal angle (θ_p) = $\theta_{\pi\sigma} - \pi/2$




Nov 5, 2000

MSL, Stanford & NASA Ames

3/20

Three Examples of Surfaces




Nov 5, 2000

MSL, Stanford & NASA Ames

4/20

Example: Reaction on Surfaces

Hydrogenation of graphene sheet and Fullerenes




H on Graphite

$$E = -4.50 \text{ eV}$$

$$\theta_p = 12.95^\circ (0.226)$$

$$\theta_s = 0.0^\circ (0.0)$$



H on C_{60}

$$E = -3.18 \text{ eV}$$

$$\theta_p = 20.56^\circ (0.359)$$

$$\theta_s = ? (0.22)$$

- Chemical reactivity difference \neq Strain energy difference ?

Nov 5, 2000


MSL, Stanford & NASA Ames

5/20

External Chemical Reaction on Fullerene

Total reaction energy

1. Straining the bonds. (+ ΔE)
 2. Breaking π bonds. (+ ΔE)
 3. Reacting with external reactant (X) in given structure. (- ΔE)
 4. Global relaxations (- ΔE)
- Strain Energy
- Binding Energy
- Temporally ignored




Nov 5, 2000

MSL, Stanford & NASA Ames

6/20

Strain Energy Contribution to Reaction

- Fix all the carbon atoms, pull only one carbon up and calculate the energy difference



1. Elastic Strain Energy

2. Independent of External Reactant

3. θ_s shift

4. θ_s increase \Rightarrow stiffer

Nov 5, 2000

MSL, Stanford & NASA Ames

7/20

Need a new title for this slide

- The relation with the pyramidal angle (θ_p)

$$|\pi\rangle = \sqrt{2} \tan(\theta_p) |s\rangle + \sqrt{1 - 2 \tan^2(\theta_p)} |p\rangle$$

- Binding energy with external reactant X

$$E_{\text{binding}} = \sqrt{2} \tan(\theta_p) E_{\text{ss}} + \sqrt{1 - 2 \tan^2(\theta_p)} E_{\text{ps}} + E_{\text{breaking}}(\theta_p)$$

$$\text{where } E_{\text{ss}} = \langle X|H|s\rangle \text{ and } E_{\text{ps}} = \langle X|H|p\rangle$$

- E_{ss} and E_{ps} depends only on external reactant X.


Nov 5, 2000

MSL, Stanford & NASA Ames

8/20

Binding Energy Dependence on Pyramidal Angle

- Hydrogen is used as the point probe.



- $E_{\text{ex}} = -5.34 \text{ eV}$
 $E_{\text{ex}} = -1.71 \text{ eV}$
- $\Delta E_{\text{straining}}$
 $-0.1 \text{ eV} \sim -0.3 \text{ eV}$
- Independent of θ_c
- Almost linear in small θ_p

Nov. 5, 2000

MSL, Stanford & NASA Ames

9/20

Total Reaction Energy Estimation

- Total reaction energy = Minimum (strain energy + binding energy)
- Small Errors ($\sim 0.1 \text{ eV}$) from ignoring global relaxation

Formation Energy Comparison with Full Relaxation Results

	Pyramidal Angle (θ_p)			Total Reaction Energy (eV)	
	Initial Value	Estimated Value	Full Relaxation	Estimated Value	Full Relaxation
Graphite	0.000	0.220	0.226	-1.50	-1.63
Nanotube	0.091	0.291	0.292*	-2.29	-2.39
C ₆₀	0.202	0.373	0.359	-3.07	-3.18

a) The forces are relaxed up to the first nearest neighbors

Nov. 5, 2000

MSL, Stanford & NASA Ames

10/20

External Chemical Reactivity (Conclusions?)

- θ_p (initial pyramidal angle) $\left\{ \begin{array}{l} \text{Strain Energy curve shift } \theta_p \\ \text{Binding Energy curve shift } < -0.3 \text{ eV} \end{array} \right.$
- Binding Energy curve - almost linear
 $\theta_p = \theta_c + \theta_s - \delta$ (stiffness change)
- Total reaction energy changing by θ_p
Strain energy effect $< 0.1 \text{ eV}$
Binding energy effect $\sim \theta_p E_{\text{ex}} \text{ eV}$

Nov. 5, 2000

MSL, Stanford & NASA Ames

11/20





Internal Reaction

- Less Reactive than External Reaction \leftarrow Less Electron Density
 - Total reaction energy
 - Straining the bonds. (+ ΔE)
 - Breaking π bonds. (+ ΔE)
 - Reacting with external reactant (X) in given structure. (- ΔE)
 - Global relaxations (- ΔE)
 - Weak bonds with neighbor carbon atoms. (- ΔE)
- Same as External Chemical Reaction
- Difference from External Reaction

Nov. 5, 2000

MSL, Stanford & NASA Ames

12/20



Quantum Computer

- Solid state quantum computer can be designed by ^{31}P atoms in bulk Si [1].



- Remaining Problems
 - Placing of a ^{31}P atom
 - Diffusion of a ^{31}P atom

- ^{31}P @ Bucky Onion
 - Possible self-ordering
 - Preventing diffusion of ^{31}P


1. Kane, B.E., *Nature*, 393, p.133 (1998)

Nov. 5, 2000

MSI, Stanford & NASA Ames

17/20

P @ the Center of C_{60}



- Full Relaxation

- Binding Energy
Center : -0.99 eV
(6,6) Bond : -0.81 eV

- Diffusion Barrier
Center \Rightarrow Bond
-0.33 eV



Nov. 5, 2000

MSI, Stanford & NASA Ames

18/20

H @ $\text{C}_{20}\text{D}_{20}$

- No meta-stable site and Diffusion Barrier = 1.17 eV



Nov. 5, 2000

MSI, Stanford & NASA Ames

19/20

Conclusions

- The external reaction energy is determined by a competition of strain energy and binding energy.
- We can estimate external reaction energy up to 0.1 eV error, which is from global relaxation.
- External Chemical Reactivity is enhanced by binding energy difference due to initial pyramidal angle.
- Inner Chemical Reactivity is enhanced by Global Interactions.
- Possible candidates for Quantum Computer
 ^{31}P @ C_{60} and ^1H @ $\text{C}_{20}\text{D}_{20}$

Nov. 5, 2000

MSI, Stanford & NASA Ames

20/20

Future Work

- We are going to test more examples on external chemical reactivity.
- We will continue to work on the reactivity of the multi-bonds reactions and non-covalent bond reactions.
- We need more systematic study on internal chemical reactivity.
- Since we found the way to encapsulate P and H, we will work on self-ordering of endo-fullerene for quantum computer application.
- As an another candidate for quantum computer, we are looking into compressed bucky onion, which has diamond core structure.

Nov 5, 2000

MSL, Stanford & NASA Ames

21/20



National
Aeronautics and
Space
Administration

NASA Scientific and Technical Document Availability Authorization (DAA)

The DAA approval process applies to all forms of published NASA Scientific and Technical Information (STI), whether disseminated in print or electronically. It is to be initiated by the responsible NASA Project Officer, Technical Monitor, author, or other appropriate NASA official for all presentations, reports, papers, and proceedings that contain NASA STI. Explanations are on the back of this form and are presented in greater detail in NPG 2200.2, "Guidelines for Documentation, Approval, and Dissemination of NASA Scientific and Technical Information."

☒ Original
☐ Modified

I. DOCUMENT/PROJECT IDENTIFICATION

TITLE

Curvature Dependent Reactivity of Fullerenes and Nanotubes

AUTHOR(S)

Srivastava, Deepak (650) 604-3486
(CSC)

S. Park and K. Cho
Stanford University

ORIGINATING NASA ORGANIZATION
NASA Ames Research Center, Code IN

PERFORMING ORGANIZATION (If different)
CSC

CONTRACT/GRANT/INTERAGENCY/PROJECT NUMBER(S)
DTTS59-99-D-00437 / A61812D

DOCUMENT NUMBER(S)

DOCUMENT DATE

For presentations, documents, or other STI to be externally published (including through electronic media), enter appropriate information on the intended publication such as name, place, and date of conference, periodical, or journal name, or book title and publisher in the next box. These documents must be routed to the NASA Headquarters or Center Export Control Administrator for approval (see Sections III and VIII).

A contributed talk at the 8th Foresight Conference on Molecular Nanotechnology, Nov 2-5, Bethesda, MD.

II. SECURITY CLASSIFICATION

CHECK ONE (One of the five boxes denoting Security Classification must be checked.)

☐ SECRET ☐ SECRET RD ☐ CONFIDENTIAL ☐ CONFIDENTIAL RD ☒ UNCLASSIFIED

III. AVAILABILITY CATEGORY

☐ ITAR ☐ EAR

Export Controlled Document - USML Category _____ /CCL Export Control
Classification Number (ECCN) _____ (Documents marked in this block must have the
concurrence/approval of the NASA Headquarters or Center Export Control Administrator (see Section VIII).)

☐ TRADE SECRET

☐ U.S. Government agencies and U.S. Government agency contractors only

☐ SBIR

☐ NASA contractors and U.S. Government only

☐ COPYRIGHTED

☐ U.S. Government agencies only

☐ NASA personnel and NASA contractors only

☐ NASA personnel only

☐ Available only with the approval of the issuing office: _____

☐ Limited until (date) _____

☒ PUBLICLY
AVAILABLE

Publicly available documents must be unclassified, may not be export controlled, may not contain trade secret or confidential commercial data, and should have cleared any applicable patents application process.

IV. DOCUMENT DISCLOSING AN INVENTION

THIS DOCUMENT MAY BE RELEASED ON (date)

NASA HEADQUARTERS, CENTER, OR INTELLECTUAL PROPERTY COUNSEL SIGNATURE

DATE

ALREADY RELEASED

[Signature]

11-21-00

V. BLANKET RELEASE (OPTIONAL)

☐ All documents issued under the following contract/grant/project number _____
may be processed as checked in Sections II and III.

☐ The blanket release authorization granted on (date) _____

☐ is RESCINDED - Future documents must have individual availability authorizations.

☐ is MODIFIED - Limitations for all documents processed in the STI system under the blanket release should be changed to conform to blocks as checked in Sections II and III.

DEC 07 2000